



Complete mitochondrial genome of the Oriental Hornet, *Vespa orientalis* F. (Hymenoptera: Vespidae)

Haddad, Nizar Jamal; Al-Nakeeb, Kosai Ali Ahmed; Petersen, Bent; Dalén, Love; Sorgenfrei Blom, Nikolaj; Sicheritz-Pontén, Thomas

Published in:
Mitochondrial DNA Part B

Link to article, DOI:
[10.1080/23802359.2017.1292480](https://doi.org/10.1080/23802359.2017.1292480)

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Haddad, N. J., Al-Nakeeb, K. A. A., Petersen, B., Dalén, L., Sorgenfrei Blom, N., & Sicheritz-Pontén, T. (2017). Complete mitochondrial genome of the Oriental Hornet, *Vespa orientalis* F. (Hymenoptera: Vespidae). *Mitochondrial DNA Part B*, 2(1), 139-140. <https://doi.org/10.1080/23802359.2017.1292480>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <http://www.tandfonline.com/loi/tmdn20>

Complete mitochondrial genome of the Oriental Hornet, *Vespa orientalis* F. (Hymenoptera: Vespidae)

Nizar Jamal Haddad, Kosai Al-Nakeeb, Bent Petersen, Love Dalén, Nikolaj Blom & Thomas Sicheritz-Pontén

To cite this article: Nizar Jamal Haddad, Kosai Al-Nakeeb, Bent Petersen, Love Dalén, Nikolaj Blom & Thomas Sicheritz-Pontén (2017) Complete mitochondrial genome of the Oriental Hornet, *Vespa orientalis* F. (Hymenoptera: Vespidae), Mitochondrial DNA Part B, 2:1, 139-140, DOI: [10.1080/23802359.2017.1292480](https://doi.org/10.1080/23802359.2017.1292480)

To link to this article: <http://dx.doi.org/10.1080/23802359.2017.1292480>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 01 Mar 2017.



Submit your article to this journal [↗](#)



Article views: 131



View related articles [↗](#)









View Crossmark data [↗](#)

Full Terms & Conditions of access and use can be found at
<http://www.tandfonline.com/action/journalInformation?journalCode=tmdn20>

MITOGENOME ANNOUNCEMENT



Complete mitochondrial genome of the Oriental Hornet, *Vespa orientalis* F. (Hymenoptera: Vespidae)

Nizar Jamal Haddad^a , Kosai Al-Nakeeb^b , Bent Petersen^b , Love Dalén^c , Nikolaj Blom^b  and Thomas Sicheritz-Pontén^b 

^aBee Research Department, National Center for Agricultural Research and Extension, Amman, Jordan; ^bDepartment of Bio and Health Informatics, Technical University of Denmark, Lyngby, Denmark; ^cDepartment of Biodiversity Informatics and Genetics, Swedish Museum of Natural History, Stockholm, Sweden

ABSTRACT

The Oriental Hornet (*Vespa orientalis*) is a social insect belonging to the Vespidae family (Wasps, Hornets, Yellowjackets), genus *Vespa* (true Hornets). The oriental hornet is a scavenger and an agricultural pest, especially to bee farmers, but is also recently described as a harvester of solar energy. Here, we report the mitochondrial genome sequence of the Oriental Hornet, *Vespa orientalis* F., which may play a vital role in understanding this wasp biology, light trapping and generation of electricity. The mitochondrial genome of this hornet is 16,099 bp in length, containing 13 protein-coding genes, 21 transfer RNA genes, and 2 ribosomal RNA genes. The overall base composition of the heavy-strand is 40.3% A, 5.9% C, 13.2% G, and 40.6% T, the percentages of A and T being higher than that of G and C. The mitochondrial genome of the Oriental Hornet, *Vespa orientalis* F. represents the first mitogenome of a solar energy harvesting insect.

ARTICLE HISTORY

Received 19 January 2017
Accepted 5 February 2017

KEYWORDS

Oriental hornet; *Vespa orientalis* F.; mitochondrial genome

The Oriental Hornet, *Vespa orientalis* F., is distributed throughout the Levant region, Southern Europe, Northeast Africa, and Southwestern Asia including India (Bodenheimer 1951; Haddad Nizar Fuchs et al. 2005). It is known to prey on various insect species, but shows a marked preference for honeybees and is thus considered to be an apicultural pest (Haddad et al. 2007). The hornet causes damage by destroying bee hives and by reducing or even inhibiting the flights of bees (Blum 1956), which results in considerable loss for beekeepers and reduced pollination of crops. It has also become a serious pest on different fruit trees (grape, date palm, pear, peach, citrus, etc.) either by directly destroying the fruits or by the indirect damage caused by saprophytic fungi. In addition, hornet stings can cause medical problems where individuals react differently to being stung: some are scarcely affected while others suffer considerable pain and swelling (Bodenheimer 1951; Haddad et al. 2007). In recent years, it was discovered that pigments in the hornet's yellow tissues trap light, while its brown tissues generate electricity. Exactly how the hornets use this electricity is still not entirely understood, but the harvested energy might be used in physical activities, like flight and in temperature regulation (Kaplan 2010; Plotkin et al. 2010). In 2012 we collected and sequenced the genome of an Oriental hornet at the Maru

field station in Amman, Jordan (32°36'22.9"N, 35°54'04.9"E). The specimen was disposed during sequencing. We assembled the genome and extracted the mitochondrial sequence from the assembly and annotated it using the MITOS webserver. The mitochondrial sequence reported here adds a new genetic resource for the family Vespidae and will contribute to the understanding of the evolution of sun light harvesting social insects.

The mitochondrial genome of *V. orientalis* is 16,099 bp in length and contains 21 tRNA genes, 13 protein-coding genes, 2 rRNA genes, and 1 non-coding control region (D-loop). The small and large rRNA subunits are 763 and 1378 bp in length, respectively. The overall base composition of the heavy-strand is 40.3% A, 5.9% C, 13.2% G, and 40.6% T. The Cytochrome B gene was split into two parts during annotation. This could be due to internal duplications in that gene; however, the total length corresponds to mitochondrial Cytochrome B genes in similar organisms. We did not observe any overlaps between genes on the same strand. The longest intergenic gap was between the 12s RNA and the gene coding for tRNA-Tyr. The intergenic gap is defined in Figure 1 as the difference between the start and end positions between two adjacent genes. The sequence is available in NCBI Genbank under accession KY563657.

CONTACT Thomas Sicheritz-Pontén  thomas@cbs.dtu.dk  Building 208, 2800 Kgs. Lyngby, Denmark

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Phylogenetic tree of the Oriental hornet with the Giant Asian hornet (*Vespa mandarinia*), Australian hornet (*Abispa ephippium*), Argentine ant (*Linepithema humile*) and the common honey bee (*Apis mellifera ligustica*) as the outgroup. The tree was created using the parsimony criterion on the ungapped sequences of a multiple sequence alignment.

Acknowledgements

The authors acknowledge support from Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the National Genomics Infrastructure (NGI) for assistance with massively parallel sequencing.

Disclosure statement

The author reports no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Funding

The authors acknowledge support from Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the National Genomics Infrastructure (NGI) for assistance with massively parallel sequencing.

ORCID

Nizar Jamal Haddad <http://orcid.org/0000-0003-0250-5291>
 Kosai Al-Nakeeb <http://orcid.org/0000-0003-3432-3628>
 Bent Petersen <http://orcid.org/0000-0002-2472-8317>

Love Dalén <http://orcid.org/0000-0001-8270-7613>
 Nikolaj Blom <http://orcid.org/0000-0001-8012-6380>
 Thomas Sicheritz-Pontén <http://orcid.org/0000-0001-6615-1141>

References

- Blum R. 1956. Bienen im Nahen Osten. Proc. Inter. Apic. Cong. (Apimondia). 16:28.
- Bodenheimer FS. 1951. Citrus entomology in the Middle East. In: W. Junk, editor. Netherlands: The Hague; p. 76–78.
- Haddad N, Dvorak L, Adwan O, Mdanat H, Bataynah A. 2007. A new data on Vespidae wasp fauna of Jordan (H. Vespidae). Linzer Biol Beitr. 39:137–142.
- Haddad Nizar Fuchs S, Haddaden J, Kopelke JP. 2005. Record of Sphecophaga vesparum Curtis, a natural enemy of the *Vespa orientalis* in the northern part of Jordan. Zool Middle East. 35:114–116.
- Plotkin M, Hod I, Zaban A, Boden SA, Bagnall DM, Galushko D, Bergman DJ. 2010. Solar energy harvesting in the epicuticle of the oriental hornet (*Vespa orientalis*). Naturwissenschaften. 97:1067–1076.
- Kaplan M. 2010. Solar-powered hornet found; turns light into electricity. National Geographics. 1–3.